

**DETERMINATION OF SCAVENGE ACTIVITY AND PHENOLIC
CONTENTS IN AQUEOUS AND ETHANOLIC EXTRACT OF
DRAGON FRUIT**

AZRUL FAHMI BIN SULAIMAN

**Final Year Project Report Submitted in
Partial Fulfillment of the Requirement for the
Degree of Bachelor of Science (Hons.) Applied Chemistry
in the Faculty of Applied Sciences
Universiti Teknologi Mara**

MAY 2009

This Final Year Project entitled **“Determination of Scavenge Activity and Phenolic Contents in Aqueous and Ethanolic Extract of Dragon Fruit”** was submitted by Azrul Fahmi Bin Sulaiman, in partial fulfillment of the requirement for the Degree of Bachelor of Science (Hons.) Applied Chemistry, in the Faculty of Applied Science, and was approved by:



Pn. Haliza bt Kassim

Supervisor


B. Sc. (Hons.) Applied Chemistry

Faculty of Applied Science

Universiti Teknologi MARA

40450 Shah Alam

Selangor



Cik Sabrina bte M yahaya

Project coordinator

B. Sc. (Hons.) Applied Chemistry

Faculty of Applied Science

Universiti Teknologi MARA

40450 Shah Alam

Selangor



Dr. Yusairie bin Mohd

Head of Programme

B. sc. (Hons) Applied chemistry

Faculty of Applied Science

Universiti Teknologi MARA

40450 Shah Alam

Selangor

Date: 28 MAY 2009

ACKNOWLEDGEMENTS

First of all, Alhamdulillah, with the blessing from ALLAH S.W.T, I am successfully to completing this final year project.

I would like to give my sincere gratitude to my dearest supervisor, Pn.Haliza binti Kassim who had really been supporting, motivating and guiding me during the process of finishing this final year project.

I would like also to extend my special thank to all staff that are involved for their help, assistant and guidance during completing this project.

Finally, I would like to thanks all my friends that had been supporting, helping, sharing and giving good ideas during the process of finishing this project proposal. May Allah bless all of you. Thanks you.

Azrul Fahmi.

TABLE OF CONTENTS

| | Page |
|---|------|
| ACKNOWLEDGEMENT | iii |
| TABLES OF CONTENTS | iv |
| LIST OF TABLES | vi |
| LIST OF FIGURES | vii |
| LIST OF ABBREVIATIONS | viii |
| ABSTRACT | ix |
| ABSTRAK | x |
| | |
| CHAPTER 1 INTRODUCTION | |
| 1.1 Background and problem statement | 1 |
| 1.2 Significance of study | 3 |
| 1.3 Objectives | 3 |
| | |
| CHAPTER 2 LITERATURE REVIEW | |
| 2.1 Dragon fruit | 4 |
| 2.2 antioxidant | 7 |
| 2.2.1 Natural antioxidant | 8 |
| 2.2.2 Synthetic antioxidant | 8 |
| 2.3 free radical | 9 |
| 2.3.1 Mechanism of free radical | 9 |
| 2.3.2 Sources of free radical | 10 |
| 2.4 total phenolic content | 11 |
| | |
| CHAPTER 3 METHODOLOGY | |
| 3.1 Materials | 12 |
| 3.1.1 Raw materials | 12 |
| 3.1.2 Chemical | 12 |
| 3.1.4 Instruments | 12 |
| 3.2 Methods | |
| 3.2.1 Extraction | |
| 3.2.1.1 Extraction with ethanol | 13 |
| 3.2.1.2 Extraction with water | 13 |

ABSTRACT

This study was carried out to determine the total phenolic content and scavenge activity as well as to determine which solvent extraction will producing high content of total phenolic content in Dragon Fruit. Two solvent extractions were used in these studies which are water and ethanol. In total phenolic method, Gallic acid was used as the standards thus the total phenolic content was expressed as mg GAE / g fresh sample. The higher total phenolic content is shown by red-ethanolic extract (1.395 ± 0.022) mg/g followed by red-aqueous extract (1.130 ± 0.029) mg/g, white-aqueous extract (0.733 ± 0.022) mg/g and the lowest is shown by white-ethanolic extract (0.569 ± 0.017) mg/g. The percentage of scavenge activity of red-aqueous extract shows the highest with 95.51%, followed by red-ethanolic extract (93.02%), white-ethanolic extract (81.38%) and the lowest is white-aqueous extract (75.81%). The overall results indicated that the red dragon fruit contain high total phenolic content and higher percentage of scavenge activity compare to white one. So, the red one is concluded to contain more antioxidant properties compare to the white one.